

Mach–Zehnder-like interferometry with graphene nanoribbon networks

Sofia Sanz^{*}, Nick Papior[†], Géza Giedke^{‡,§}, Daniel Sánchez-Portal[¶],
Mads Brandbyge^{*}, Thomas Frederiksen^{‡,§}

^{*}Department of Physics, DTU, DK-2800 Kgs. Lyngby, Denmark

[†]DTU Computing Center DK-2800 Kgs. Lyngby, Denmark

[‡]Donostia International Physics Center (DIPC), E-20018, Donostia-San Sebastián, Spain

[§]IKERBASQUE, Basque Foundation for Science, E-48013, Bilbao, Spain

[¶]Centro de Física de Materiales CSIC-UPV/EHU, E-20018, Donostia-San Sebastián, Spain
e-mail: sofsa@dtu.dk

INTRODUCTION

The similarities between electrons traveling ballistically in graphene nanostructures and photons propagating in waveguides has spawned the novel field of electron quantum optics in graphene constrictions. With the ideas of optics-like experiments in mind, we can think of performing electron interferometry in graphene constrictions, for which some elementary building blocks, e.g., such as electron beam splitters, mirrors, etc., are needed. Remarkably, it has been recently shown that electrons injected in devices formed of two infinite graphene nanoribbons (GNR) placed one on top of the other with a crossing angle of 60° can be split into two outgoing waves without reflection [1], [2], [3]. Moreover, GNRs with zigzag edge topology are expected to host spin polarized states due to magnetic instabilities of the localized states at the edges [4], which make these devices even more interesting since we can think of performing both electron and spin quantum optics experiments [5]. With the advent of on-surface synthesis techniques, now not only atomistic defect-free samples of GNRs can be produced [6], [7] but also they can be manipulated.

In this work we propose interesting networks for studying electron and spin quantum interferometry built from four crossed GNRs in a pairwise setup (see Figure 1) that splits the beam into two possible paths that will self-interfere at the outgoing ports, where, the resulting interference pattern can be further tuned by an external magnetic field [8] as a




Fig. 1. Geometry of GNR-based interferometer with the two interfering paths sketched.

consequence of the Aharonov-Bohm effect [9]. By means of the mean-field Hubbard model [10], [11] in combination with the non-equilibrium Green's function formalism (NEGF) we are able to describe the spin dependent transport in this multi-terminal device. We further show that the scattering matrix formalism in the approximation of independent scattering at the four individual junctions provides accurate results as compared with the Green's function description, allowing for a simple interpretation of the interference process between two dominant pathways.

REFERENCES

- [1] L. R. F. Lima, A. R. Hernandez, F. A. Pinheiro, and C. Lewenkopf, A 50/50 electronic beam splitter in graphene nanoribbons as a building block for electron optics, *J. Phys.: Condens. Matter* **28**, 505303 (2016).
- [2] P. Brandimarte, M. Engelund, N. Papior, A. Garcia-Lekue, T. Frederiksen, and D. Sanchez-Portal, A tunable electronic beam splitter realized with crossed graphene nanoribbons, *J. Chem. Phys.* **146**, 092318 (2017).

- [3] S. Sanz, P. Brandimarte, G. Giedke, D. Sanchez-Portal, and T. Frederiksen, Crossed graphene nanoribbons as beam splitters and mirrors for electron quantum optics, *Phys. Rev. B* **102**, 035436 (2020).
- [4] Y. W. Son, M. L. Cohen, and S. G. Louie, Half-metallic graphene nanoribbons, *Nature* **444**, 347 (2006).
- [5] S. Sanz, N. Papior, G. Giedke, D. Sanchez-Portal, M. Brandbyge, and T. Frederiksen, Spin-polarizing electron beam splitter from crossed graphene nanoribbons, *Phys. Rev. Lett.* **129**, 037701 (2022).
- [6] J. Cai, P. Ruffieux, R. Jaafar, M. Bieri, T. Braun, S. Blankenburg, M. Muoth, A. P. Seitsonen, M. Saleh, X. Feng, K. Mullen, and R. Fasel, Atomically precise bottom-up fabrication of graphene nanoribbons, *Nature* **466**, 470 (2010).
- [7] P. Ruffieux, S. Wang, B. Yang, C. Sanchez-Sanchez, J. Liu, T. Dienel, L. Talirz, P. Shinde, C. A. Pignedoli, D. Passerone, T. Dumslaff, X. Feng, K. Mullen, and R. Fasel, On-surface synthesis of graphene nanoribbons with zigzag edge topology, *Nature* **531**, 489 (2016).
- [8] S. Sanz, N. Papior, G. Giedke, D. Sanchez-Portal, M. Brandbyge, and T. Frederiksen, Mach-Zehnder-like interferometry with graphene nanoribbon networks, arxiv.org/abs/2302.04821.
- [9] Y. Aharonov and D. Bohm, Significance of Electromagnetic Potentials in the Quantum Theory, *Phys. Rev.* **115**, 485 (1959).
- [10] J. Hubbard, Electron correlations in narrow energy bands, *Proc. R. Soc. A* **276**, 238 (1963).
- [11] S. S. Wuhl, N. Papior, M. Brandbyge, and T. Frederiksen, hubbard: v0.2.0, doi.org/10.5281/zenodo.4748765.